



Case report

Toxicology observation: Nystagmus after marijuana use

Thomas Kibby MD, MPH Instructor*, S. Eliza Halcomb MD Assistant Professor

Toxicology Section, Emergency Medicine, Washington University School of Medicine, St. Louis, MO, USA

ARTICLE INFO

Article history:

Received 10 November 2011

Received in revised form

28 April 2012

Accepted 28 July 2012

Available online 1 September 2012

Keywords:

Forensic science

Marijuana abuse/detection

Humans

Psychomotor performance/drug effects

Nystagmus

Pathologic/aetiology

Substance abuse detection

ABSTRACT

Traditional teaching has held that horizontal-gaze nystagmus is a sign of intoxication by sedatives such as alcohol but not marijuana. This is a case report of an adult male who presents with 3 days of visual disturbance and dizziness following marijuana use. The exam was notable for gaze-evoked nystagmus and ataxia. Lab testing was normal except that urine drug screening was positive for marijuana only. Imaging included computed tomography (CT) and magnetic resonance imaging (MRI) scans of the head. Prior studies showing a negative association of nystagmus with marijuana are reviewed. This case is presented as a possible exception to the generalisation that marijuana is not associated with nystagmus.

© 2012 Elsevier Ltd and Faculty of Forensic and Legal Medicine. All rights reserved.

1. Introduction

The Standardized Field Sobriety Tests (SFSTs)¹ are widely used by non-clinical personnel to detect clinical intoxication. They are typically used as a pre-arrest screen by law enforcement personnel, and have been validated specifically for impairment by ethanol.² As part of the exam, the tester directs the subject into lateral and upward gaze, observing for smooth pursuit tracking and nystagmus. The presence of nystagmus to lateral or vertical gaze or onset of less than 45° deviation is considered suggestive of intoxication. Traditional training for the SFST indicates that the presence of nystagmus at various points is suggestive of intoxication by sedatives including alcohol, barbiturates and phencyclidine (PCP), but not marijuana. The following is a description of a case in which marijuana use is associated with nystagmus.

2. Materials and methods: case report

The patient is a 25-year-old male who presented to a local emergency department with a 3-day history of persistent visual disturbance and dizziness. He noted that his symptoms were aggravated by gazing laterally or with change in head position. He

also noted difficulty in coordination during daily activities such as dressing and walking. He had no prior medical or surgical problems; he was not currently using medications. His review of systems was positive for nausea, vomiting, and a limited recent headache. He denied fever, chills or tinnitus. He smoked 4 g of marijuana and drank several beers 4 days and again 3 days prior to presentation. He denied ethanol or marijuana consumption for the 2 days before presentation. He denied focal weakness or numbness, dysarthria, aphasia or other symptoms.

On exam the patient was awake, alert and appropriate. His vital signs were normal. Pertinent positive findings by neurologist's exam were left end-gaze sustained nystagmus, rotational nystagmus to vertical gaze and ataxia. The remainder of the neurologic exam was normal. Ear, nose and throat (ENT), cardiac, pulmonary and abdominal exams were normal.

All of his laboratory work was normal except for the urine drug screen, which was positive for cannabinoids and negative for PCP, amphetamines, barbiturates, benzodiazepines, cocaine and opiates. A biologic fluid screen for ethanol was not obtained. A computed tomography (CT) scan of the head and magnetic resonance imaging (MRI) scan of the head, cerebellum and brainstem were both negative for pathology. The electrocardiogram showed normal sinus rhythm with normal intervals. His symptoms markedly improved after administration of intramuscular (IM) lorazepam, to the point of becoming symptom-free and non-ataxic. Both neurology and toxicology consultants concluded that the vertigo and nystagmus were due to marijuana use. The patient was advised

* Corresponding author. 660 S. Euclid, CB 8072, St. Louis, MO 63110, USA. Tel.: +1 314 747 5800; fax: +1 314 747 5866.

E-mail address: kibbyt@wustl.edu (T. Kibby).

to discontinue its use, and was released with directions to follow with his personal physician.

3. Discussion

The oculomotor effects of marijuana were first investigated during the period when such effects by alcohol were described. Spector³ described the absence of nystagmus as measured by electronystagmography recordings in 72 volunteers who smoked marijuana. Flom⁴ and Baloh⁵ described the oculomotor effects of alcohol including inhibition of smooth tracking and saccadic tracking, but neither found significant effect when marijuana was studied alone and the latter found marginal additive effect when combined with alcohol. These studies supported the conclusion that marijuana use does not cause nystagmus on the SFST. A later study⁶ found impairment of smooth pursuit eye tracking with marijuana, especially at higher doses and tracking in lateral fields, but neither nystagmus nor saccadic tracking were studied. One recent study⁷ found that test subjects also displayed a lack of smooth pursuit but not nystagmus after smoking marijuana. Dizziness is a well-described adverse effect of marijuana, especially in studies of the medical use of cannabinoids.⁸ However, it is unusual for vertiginous symptoms to persist for several days. There have been a number of case reports of stroke or transient ischaemic attacks (TIA's) associated with heavy marijuana use, recently summarised by Thanvi and Treadwell.⁹ Of the nine case series reviewed with imaging studies, eight reported positive studies for infarcts and the ninth involving TIA's reported only white-matter changes. Of interest is that the majority of the infarcts reported in this review involved the cerebellum.

Our patient had onset of dizziness and visual changes after consuming marijuana and alcohol, which persisted 2 days later. In retrospect, it would have been useful to know if other adulterants were present but the source marijuana and specific testing were not available. The association of this patient's nystagmus with

marijuana would have been stronger had testing been performed to exclude the presence of ethanol. Although there are anecdotal reports of vertigo after marijuana use on the Internet, there are only two published studies relating the oculomotor effects of cannabinoid use and in these nystagmus has not been described. This case is presented as a possible exception to the generalisation that marijuana is not associated with nystagmus.

Ethical approval

None.

Funding

No financial support is reported.

Conflict of interest

None.

References

1. Stuster J. *Development of a standardized field sobriety test (SFST) training management system*. National Highway Transportation Safety Administration; 2001. DOT HS 809 400, Appendix A.
2. Stutser J. Validation of the standardized field sobriety test battery at 0.08% blood alcohol concentration. *Hum Factors* 2006;**48**:608–14.
3. Spector M. Acute vestibular effects of marijuana. *J Clin Pharmacol* 1973;**13**:214–7.
4. Flom MC, Brown B, Adams AJ, Jones RT. Alcohol and marijuana effects on ocular tracking. *Am J Optom Physiol Opt*. 1976;**53**:764–73.
5. Baloh RW, Sharma S, Moskowitz H, Griffith R. Effect of alcohol and marijuana on eye movements. *Aviat Space Environ Med*. 1979;**50**:18–23.
6. Fant RF, Heishman SJ, Bunker EB, Pickworth WB. Acute and residual effects of marijuana in humans. *Pharmacol Biochem Behav*. 1998;**60**:777–84.
7. Papafotiou K, Carter JD, Stough C. An evaluation of the sensitivity of the standardised field sobriety tests (SFSTs) to detect impairment due to marijuana intoxication. *Psychopharmacol* 2005;**180**:107–14.
8. Wang T, Collet J, Shapiro S, Ware MA. Adverse effects of medical cannabinoids: a systematic review. *CMAJ* 2008;**178**:1669–78.
9. Thanvi BR, Treadwell SD. Cannabis and stroke: is there a link? *Postgrad Med J* 2009;**85**:80–3.